

Bactericidal and fungicidal liquid preparations for  
industrial products

The present invention relates to compositions  
5 for use in industrial products which protect these  
products against bacterial and fungal attack over  
relatively long service lives. These compositions are,  
in particular, themselves stable and comprise at least  
one bactericidal N-formal, at least one fungicide and  
10 at least one stabilizer. The invention further relates  
to industrial products to which these compositions have  
been added.

Preservatives with microbicidal action for use  
in industrial products such as cooling lubricants,  
15 cooling lubricants mixed with water, industrial-grade  
emulsions and other water-based industrial products,  
but also in household products, such as, for example,  
cleaning products or cosmetics, such as, for example,  
bodycare compositions, disinfectants and seed treatment  
20 compositions are generally added to the products to be  
preserved as required in the form of concentrates in a  
small amount.

These preservatives protect these products  
against bacterial and fungal attack and contribute to  
25 long service lives and long usability and applicability  
of industrial products, and household and cosmetic  
products.

During their manufacture, storage and their  
use, the preservatives are subject to certain  
30 requirements which arise, inter alia, from the way in  
which they are added to said products in the form of  
concentrates. These concentrates should themselves be  
sufficiently stable over long service lives.

Known bactericidal active ingredients which are  
35 currently frequently used are N-formals which, in  
addition to the biocidal, in particular bactericidal,  
action, have a (desired) alkalinizing and buffer  
action. N-formals are reaction products of aldehydes  
with amines. These N-formals act as aldehyde donor

compounds. In order to achieve microbicidal broadband action, it is, however, necessary to improve the fungicidal action of the N-formal-containing compositions or industrial products. However, 5 incompatibilities are frequently found between N-formals and fungicides, which are evident from the decrease in active ingredient content, resulting in inadequate effectiveness. These problems arise irrespective of whether the components N-formal and 10 fungicide are added to the industrial products simultaneously or separately, i.e. during storage of the preservative and in the product treated therewith.

Ways of improving the stability of compositions which comprise bactericidal N-formals and fungicides 15 have therefore been sought.

DE 197 05 085 A1 describes algicidal and fungicidal preservatives which comprise a pyrrithione, an algicidal triazine and a fungicidal derivative of benzimidazole or thiophene.

20 GB 2 274 779 A describes dilute fungicidal compositions for use in aerosols which consist of an aromatic sulphur-containing compound or a mixture of an aromatic sulphur-containing compound and a triazine. In the comparative examples, the disadvantages of these 25 compositions are compared with the additive mixtures according to the invention.

DE 197 22 858 discloses compositions with bactericidal and fungicidal action, which comprise a formaldehyde donor compound and an iodopropynyl 30 compound and may further comprise isothiazolones or 2-mercaptopyridine N-oxide.

DE 195 34 532 C2 discloses, as stabilizers for isothiazolones, 2-mercaptopyridine N-oxide and corresponding salts and mild oxidizing agents, such as, 35 for example, hydrogen peroxide or t-butyl hydroperoxide.

The attempt to formulate ready-to-use preparations according to the prior art have however led to inhomogeneous or unstable solutions, even after

short storage times. These inhomogeneities hinder exact dosing of the composition or make it impossible, this being true in particular for N-formal compositions. Although a separate addition of active ingredient and stabilizer is possible, in practice it is not desired and is plant- and labour-intensive, and is susceptible to errors.

On the other hand, in compositions which necessarily prescribe the presence of more than three components as primary active ingredients, the possibility of being able to mix within broad concentration ranges is limited, and the costs are also considerably higher.

Accordingly, the object of the present invention is to make available additive mixtures for cooling lubricant products which comprise N-formals as bactericidal active ingredients and also fungicides, which protect the industrial products from microbial attack and improve the shelf life and service life of the industrial products. The compositions according to the invention should in themselves be sufficiently stable over a long period and be storable and not decompose under practical conditions.

A further object of the present invention is to formulate compositions comprising N-formal and fungicide such that they can be metered easily to commercially available industrial products, e.g. by adding a liquid preparation. It is also the object of the present invention to provide effective industrial products which are characterized in that they have an increased service life and improved effectiveness compared with the prior art.

A further object was to develop compositions which necessarily prescribe the presence of the fewest possible different active ingredient components, the intention being for the long term stability and the high effectiveness to be retained.

These objects are achieved by the characterizing features of Patent Claim 1. According to

Patent Claim 1, the stable microbicidal composition is characterized in that it comprises at least one bactericidal N-formal, at least one fungicide and at least one stabilizer.

5 In addition to the customary constituents, the industrial products comprise an additive mixture according to the invention. The additive mixture is added to the industrial products, in particular crop-protection compositions, seed treatment compositions,  
10 industrial preservatives, in particular pack preservatives, cooling lubricant additives, fuel additives, disinfectants, in particular low-foam disinfectants, compositions for controlling cut wounds, parasites and plants, compositions for treating plant  
15 cut wounds, film preservatives for outside and, in particular, inside, disinfectants in areas where increased fungal attack is to be expected, and wood preservatives, in a concentration greater than 0.01% by weight, preferably greater than 0.05% by weight and in  
20 particular greater than 0.10% by weight, based on the weight of the industrial product. For example, 0.01 to 10% by weight, preferably 0.05 to 5% by weight and in particular 0.10 to 1% by weight, based on the weight of the industrial product, are added. The components of  
25 the composition can be incorporated into the industrial products separately from one another, in particular temporally separately from one another, it being possible for the components of the composition (or the composition) to be incorporated with the controlled  
30 introduction of heat.

The N-formals according to the invention are condensation products of aldehydes, preferably formaldehyde (e.g. p-formaldehyde), acetaldehyde or propionaldehyde, in particular formaldehyde and  
35 p-formaldehyde, and amines, e.g. primary or secondary amines, preferably alkanolamines and oxazolidines (e.g. monoethanolamine, isopropanolamine, 3-amino-1-propanol, 5-methyloxazolidine). The N-formal is generally present in the additive mixture according to the invention in a

concentration of from 1 to 99% by weight, preferably from 10 to 95% by weight, more preferably 20 to 90% by weight, in particular 40 to 90% by weight, e.g. 50% by weight or 90% by weight.

5            Particularly preferred N-formals are Grotan BK (2,2',2''-(hexahydro-1,3,5-triazine-1,3,5-triyl)tri-ethanol) and Mar 71 (3,3'-methylenebis(5-methyloxazolidine)).

10           The fungicides used in the additive mixture according to the invention include isothiazolones, in particular the isothiazolones known from DE 195 34 532 C2 mentioned at the beginning. Preferred examples of isothiazolones are Kathon 893 T (2-n-octyl-2H-isothiazolin-3-one), Kathon 893 (45% strength  
15           solution of 2-n-octyl-2H-isothiazolin-3-one in 1,2-propylene glycol), benzisothiazolone, Kathon 886 (14% strength aqueous solution of a 5:1 mixture of 5-chloro-2-methyl-4-isothiazolin-3-one and 2-methyl-4-isothiazolin-3-one) or Zonen-Fex (mixture of 10%  
20           5-chloro-2-methyl-4-isothiazolin-3-one, 1% of 2-methyl-4-isothiazolin-3-one, 62.3% of ethylene glycol and 26.7% of ethylene glycol diacetate).

25           The fungicide is present in the additive mixture according to the invention in a concentration of from 0.1 to 99% by weight, preferably 2 to 40% by weight, in particular 5 to 10% by weight, e.g. 8% by weight.

30           The stabilizer can likewise have microbicidal action. Suitable stabilizers or stabilizer mixtures in the additive mixture according to the invention are 2-mercaptopyridin N-oxide and corresponding salts, preferably alkali metal and ammonium salts, such as, for example, Pyrion-Na (40% strength aqueous solution of 2-mercaptopyridine N-oxide sodium salt),  
35           2-mercaptopyridine N-oxide metal salt complexes, such as zinc pyrithione (e.g. as 48% strength aqueous dispersion), 2,2'-dithiobis(pyridine N-oxide) (Pyrion disulphide), 2-mercaptobenzothiazole, 2-(thiocyanomethyl)thiobenzothiazole, NaBrO<sub>3</sub>, and mixtures thereof.

In particular, Pyrion-Na, zinc pyrithione, Pyrion disulphide,  $\text{NaBrO}_3$ , and 2-mercaptobenzothiazole are preferred, e.g. also a mixture of Pyrion-Na and  $\text{NaBrO}_3$ . The stabilizer is present in the additive mixture according to the invention in a concentration of from 0.1 to 40% by weight, preferably 2 to 20% by weight, in particular 5 to 10% by weight, e.g. 8% by weight.

In addition, solvents and/or solubility promoters may be present in the additive mixture according to the invention, e.g. alcohol, ether, glycol, glycol monoether, glycol diether, polyol, polyol monoether, completely or incompletely etherified polyol and/or mixtures thereof, preferably phenoxy-ethanol, phenoxypropanol, 1,2-propylene glycol, 1-methoxy-2-propanol, diethylene glycol butyl ether and dipropylene glycol. Preferably, liquid additive mixtures with a high active ingredient content (e.g. > 50% by weight, preferably > 65% by weight, in particular > 75% by weight) and, accordingly, a comparatively low solvent/solubility promoter content (e.g. < 50% by weight, preferably < 35% by weight, in particular < 25% by weight) are claimed. The additive mixture can also be in liquid-viscous or paste form.

Preferably, the compositions are anhydrous or at least have a low content of water.

In addition, the composition can comprise complexing agents, e.g. phosphates and polyphosphates, ethylenediaminetetraacetic acid, nitriloacetic acid, N,N-bis(2-hydroxyethyl)glycine, diethylenetriamine-pentaacetic acid, hydroxyethanediphosphonic acid, gluconic acid, hydroxyethylethylenediaminetriacetic acid, polyoxycarboxylic acid, tris(aminomethyl)-phosphonic acid, diethylenetriaminepentamethylene-phosphonic acid, ethylenediaminetetramethylene-phosphonic acid, ethylenediaminedisuccinic acid, ethylenediaminediglutamic acid, iminodisuccinic acid, polyaspartic acid and methylglycinediacetic acid, and corresponding salts of said acids or mixtures of the complexing agents.

Moreover, corrosion-protective agents, preferably phosphonobutanetricarboxylic acid and its salts, derivatives of triazole, e.g. benzotriazole and methylbenzotriazole, Irgamet 42 (2,2'-[[ (methyl-1H-benzotriazol-1-yl)methyl]imino]bisethanol), Irgamet 39 (N,N-bis(2-ethylhexyl)-4-methyl-1H-benzotriazole-1-methylamine), and carboxylic acid derivatives, e.g. DiACID 1550 (5 (or 6)-carboxy-4-hexylcyclohex-2-en-1-octanoic acid), may be present.

Further microbicidal active ingredients, in particular O-formals, additives, in particular phenols, and/or auxiliaries may optionally be added to the additive mixture. However, the additive mixture according to the invention is preferably free from iodopropynyl compound and derivative of benzimidazole or thiophene.

The additive mixtures are prepared by simple mixing, e.g. if the N-formal is initially introduced, the stabilizer (e.g. Pyrion-Na, 40% strength in water or preferably anhydrous) is dissolved with stirring and, where appropriate, the solvents and/or solubility promoters are stirred in to give a homogeneous mixture. The fungicide is then added with stirring. If necessary, slight clouding which occurs can be separated off, e.g. filtered off, e.g. within 4, 10 or 24 hours after mixing.

During the mixing or after the mixing, the concentrates are preferably subjected to a controlled heat treatment which contributes to an improvement in the stability of the additive mixtures. Thus, for example, the mixture can be heated for 5 min to 3 hours at a temperature between 40°C and 100°C, preferably for 10 min to 2 hours at a temperature between 50°C and 90°C, in particular for 20 min to 1.5 hours at a temperature between 65°C and 85°C, e.g. 1 hour at 80°C. Furthermore, the composition can be stored for several months, such as, for example, 3 months, at a temperature of from 20°C to 50°C, in particular 30°C to 40°C.

According to the invention, it has surprisingly been found that

5 a. N-formal-containing compositions can be long-term stabilized,

b. compositions which comprise N-formal and fungicides can be long-term stabilized merely by adding a single stabilizing component,

10 c. heat treating the compositions dramatically improves their long-term stability.

The additive mixture is readily dosable, and is also storable and stable under practical conditions, i.e. at 10°C to 40°C, for several months to several  
15 years. Also advantageous is the good handlability of the additive mixture, compared with the storage, preparation and metered addition of active ingredients and stabilizers present in two-component systems.

The increased stability of the additive  
20 mixtures according to the invention is evident in particular from the lower or negligible tendency for forming sediments and clouding.

The additive mixtures according to the invention can be added in an effective manner to  
25 industrial products so that the stability thereof and the service life of the finished industrial products is improved compared with known systems.

The surprising and at times synergistic effect achieved when an additive mixture according to the  
30 invention is used in industrial products is illustrated by the examples below.



**Examples**

Percentages are given in % by weight.

In the examples below the following references apply:

5

N-formal	Mar 71	3,3'-methylenebis(5-methyloxazolidine)
	Grotan BK	2,2',2''-(hexahydro-1,3,5-triazine-1,3,5-triyl)triethanol
Fungicide	Kathon 893	a 45% strength 2-octyl-2H-isothiazol-3-one solution in 1,2-propylene glycol
	Kathon 886	a 14% strength aqueous solution of a 5:1 mixture of 5-chloro-2-methyl-4-isothiazolin-3-one and 2-methyl-4-isothiazolin-3-one
Stabilizer	Pyrion-Na	a 40% strength aqueous 2-mercapto-pyridine N-oxide sodium salt solution

**Example 1**

**Increase in the stability of a mixture of Grotan BK  
N-formal and Kathon 893 fungicide by adding the  
stabilizer Pyrion-Na**

10

	A (parts by weight)	B (parts by weight)
Grotan BK anhydrous	55.65	55.65
Kathon 893	17.0	17.0
Na-Pyrion 40%	-	13.9
Total	72.65	86.55

15 In Example A, Grotan BK (anhydrous) was mixed with Kathon 893, giving a solution which, after storage for just 3 days at room temperature, turns cloudy and forms considerable sediment. In Example B, Pyrion-Na was stirred into Grotan BK, and then Kathon 893 was added. This gives a clear solution which does not form sediment even after storage for 3 months at room  
20 temperature.

After preparation and storage at 40°C, solution A is very cloudy with a sediment after just

24 h. In contrast, after preparation and storage at 40°C, solution B is still clear and free from sediment even after 3½ months.

5 Result:

An addition of Pyrion-Na stabilizer stabilizes in accordance with the invention a preparation of Grotan BK (anhydrous) N-formal and Kathon 893 fungicide.

10

**Example 2**

**Better stability of the preparations according to the invention compared with mixtures of N-formal and stabilizer or fungicide and stabilizer**

15

A 89% Mar 71 N-formal and 11% Pyrion-Na stabilizer (40% strength in water) were mixed. The immediately yellow, opaque solution is weakly opaque with a voluminous precipitate after storage for 3 weeks at room temperature.

20

B 24.5% Kathon 893 fungicide, 17.5% Pyrion-Na stabilizer (40% in water) and 58% phenoxypropanols were mixed. The initially clear, yellow-brown solution forms a crystalline precipitate (about 5%) in yellow solution after storage at room temperature after about 4 months.

25

C 46.8% Mar 71 N-formal, 17.0% Kathon 893 fungicide, 13.4% Pyrion-Na stabilizer and 22.8% 1,2-propylene glycol were heated for 1 hour at 80°C. Cooling gives a clear, yellow-brown solution. After storage for 4 months at room temperature, this solution remains clear yellow-brown.

30

Result:

The stability of a combination of the N-formal Mar 71 and stabilizer (A) is significantly increased according to the invention by the addition of the fungicide

35

Kathon 893 (C). In addition, the addition of the N-formal Mar 71 (C) increases, in accordance with the invention, the stability of a combination of Kathon 893 and the stabilizer Pyrion-Na (B).

5

### Example 3

**Long-term stability of an additive mixture according to the invention upon storage at elevated temperature**

10           90% Mar 71 N-formal were mixed with 10% of a mixture of 24.5% Kathon 893 fungicide, 17.5% Pyrion-Na stabilizer (40% strength in water) and 58% phenoxypropanols, which gave a clear, yellow solution. After storage for 24 hours at room temperature, the solution  
15 was slightly cloudy and was filtered. Various samples were stored at -5°C and 4°C in clear glass and at 25°C and 40°C in polyethylene over a relatively long period.

#### Result:

20 After 9 weeks, all samples remained clear yellow. After storage for 12 months a slight yellow precipitate in unchanged clear yellow solution, in the case of the 40°C sample significantly less precipitate than in the other samples.

25

### Example 4

**Further improvement in the stability of the preparations according to the invention by targeted short-term heat treatment**

30

A mixture (sample A) of 46.8% Mar 71, 17% Kathon 893, 13.4% Pyrion-Na (40% in water) and 22.8% 1,2-propylene glycol was obtained. A second sample (sample B) was mixed as sample A, but heated for 1 hour  
35 at 80°C. The samples A and B according to the invention were each stored at room temperature and at 40°C for 3 months.

Result:

After storage at room temperature or 40°C, sample B has in each case significantly less precipitate than sample A. Short-term (here 1 hour) heating at 80°C has  
5 a positive effect on the stability. Furthermore, storage at 40°C leads to a reduction in the slight tendency towards clouding.

**Example 5**

10       **Improvement in the stability by heat treatment  
          depending on the solvent/solubility promoter**

The experiments of Example 4 were repeated but using 1-methoxy-2-propanol instead of 1,2-propylene  
15 glycol. Samples A and B (preparation according to Example 4) were stored at room temperature or 40°C for 3 months.

Result:

20 Samples A and B confirm the result of Example 4 that short-term heat treatment of the preparations and also storage at elevated temperature have a positive effect on the stability (with regard to precipitations). Moreover, in the case of 1,2-propylene glycol  
25 (Example 4), the slight tendency towards clouding of the samples is lower compared with the 1-methoxy-2-propanol used in Example 5.

**Example 6**

30       **Particular stability of anhydrous additive mixtures**

61% of a mixture, dewatered on a rotary evaporator, of Grotan BK N-formal (69.1 parts by weight) and Pyrion-Na stabilizer, 40% strength  
35 (13.9 parts by weight) were stirred with 17% Kathon 893 fungicide and 22% solvent/solubility promoter at 45°C to give a homogeneous mixture. The solvents/solubility promoters used were phenoxyethanol, phenoxypropanols, 1-methoxy-2-propanol, dipropylene glycol or diethylene

glycol butyl ether. In all cases, clear to slightly opaque homogeneous stable preparations were obtained.

Result:

- 5 Grotan BK N-formal can be combined with Kathon 893 fungicide to give a stable preparation if Pyrion-Na stabilizer is added, water is removed and organic solvents intended therefor are added. Because of their particularly low tendency towards clouding, solutions  
10 in diethylene glycol butyl ether (in preference to dipropylene glycol and 1-methoxy-2-propanol) have proven particularly suitable.

**Example 7**

- 15 **The effectiveness of the additive mixtures according to the invention in the S&M Boko test after incorporation into cooling lubricants**

20 The three test additive mixtures below were prepared:

- A 46.8% Mar 71, 13.4% Pyrion-Na (40% strength in water), 17% Kathon 893 and 22.8% 1,2-propylene glycol were mixed and stored at 40°C for 3 months,  
25 then stored at room temperature. The effectiveness of test product A was tested 3 years after preparation.
- B Corresponds to test product A except that after mixing the constituents of the additive mixture, the  
30 sample was held at 80°C for 1 hour. It was further stored for 3 months at 40°C, then at room temperature. The effectiveness of the test product B was tested 3 years after preparation.
- C Corresponds in the composition to test products A and B, except that the batch was held at 80°C for  
35 1 hour and the effectiveness was tested using a test product freshly prepared in this way.

5 The test products A, B and C were incorporated, in the concentrations given below, into 100 ml in each case of a 4% strength dilute solution of the cooling lubricant Almasol EP or product 287/17-1 obtainable from Castrol in tap water from Norderstedt. The growth control used was an unpreserved sample in each case.

10 Two days after the incorporation of the preservatives, the test batches were infected for the first time with 1 ml of an inoculation solution. The inoculation solution is a suspension of the microbes (cultivated on nutrient media, then adapted to cooling lubricants diluted with water) and has a titre of at least  $10^7$  microbes per ml. The inoculation was carried out with a bacterial suspension (mixed suspension of 15 *Escherichia coli*, ATCC 11229, *Klebsiella pneumoniae*, ATCC 4352 and *Pseudomonas aeruginosa*, ATCC 15442), fungal suspension (*Fusarium oxysporum*, ATCC 62318) or a mixed suspension of bacteria and fungi.

20 The test batches were subsequently inoculated twice weekly and streaked out on agar plates twice weekly, the first streaking being carried out immediately after reinoculation. The microbial growth of the streaks is assessed after incubation for three 25 days at 25°C. To be on the safe side, negative streaks were observed for a further two days and reassessed. The preservative action of the individual product concentrations is assessed by reference to the presence or absence of growth of the individual streaks.

Almasol EP cooling lubricant				
Castrol mineral oil based, amine-containing				
		Endured inoculation cycle upon inoculation with:		
	Use conc.	Bacterial suspension	Fungal suspension	Mixed suspension
	0	2	0	0
A	0.20%	>10	>12	9
	0.15%	6	>12	8
	0.10%	6	>12	8
B	0.20%	>10	>12	11
	0.15%	>10	>12	9
	0.10%	>12	>12	9
C	0.20%	>12	>12	10
	0.15%	>12	>12	9
	0.10%	>11	>12	9

Product 287/17-1 cooling lubricant				
Castrol synthetic				
		Endured inoculation cycle upon inoculation with:		
	Use conc.	Bacterial suspension	Fungal suspension	Mixed suspension
	0	6	0	0
A	0.20%	>12	>12	10
	0.15%	>12	>12	9
	0.10%	>12	>12	9
B	0.20%	>12	>12	10
	0.15%	>12	>12	9
	0.10%	>11	>12	10
C	0.20%	>12	>12	10
	0.15%	>12	>12	10
	0.10%	>12	>12	10

5 The preserved cooling lubricants show no visible differences compared with the blank in each case (cooling lubricant without active ingredient).

The effectiveness of the preparations stored for 3 years and freshly prepared against bacteria and fungi is excellent. The effectiveness in the synthetic cooling lubricant is somewhat better than in the mineral oil-based cooling lubricant. Furthermore, the results show that controlled heat treatment (here 1 hour at 80°C) significantly improves the bactericidal action of the additive mixtures, and an additive mixture briefly heat-treated in this way shows virtually no loss in action even upon storage at room temperature for many years. In particular, the effectiveness is not impaired as a result of storage at 40°C for three months.

In the case of the use in mineral oil-based cooling lubricants, a comparison of Examples A and B shows the improved bactericidal effectiveness after heating the composition according to the invention at 80°C for just one hour.